

task_tx86fewvysrcy8fc_with_calculation

Student Group

First Name	Surname	Matrikel Nr.

Table of Contents

Exercise E5 Electron Velocity in Semiconductors (written test, approx. 6 % of a 120-minute
written test, SS2022) 2

electrostatic, electric field strength, exam ee2 SS2022

Exercise E5 Electron Velocity in Semiconductors (written test, approx. 6 % of a 120-minute written test, SS2022)

A current of $I = 1 \text{ mA}$ flows through a cross-sectional area $A = 10 \text{ } \mu\text{m}^2$ in a semiconductor.

The electron density in the semiconductor is given by the number of dopant atoms per volume.

The doping shall provide 1 donor atom (= one electron) per 10^{10} silicon atoms.

The molar volume of silicon is $V_{\text{mol,Si}} = 12 \text{ } \mu\text{mol}^{-3}$, with $N_{\text{A}} = 6.022 \cdot 10^{23}$ silicon atoms per 1 mol .

The elementary charge is given as: $e_0 = 1.602 \cdot 10^{-19} \text{ As}$

What is the average electron velocity v_e in this semiconductor?

Path

The following formula gives the speed, where n_e is the number of electrons per volume.

$$v_e = \frac{I}{n_e \cdot e_0 \cdot A}$$

n_e can be derived from the overall number of Si-atoms per volume ($\frac{N_{\text{A}}}{V_{\text{mol,Si}}}$) and the fraction k_{Donators} of these atoms, which got substituted by donators.

$$n_e = \frac{N_{\text{A}}}{V_{\text{mol,Si}}} \cdot k_{\text{Donators}} \cdot e_0 \cdot A$$

Putting in the numbers:

$$v_e = \frac{1 \cdot 10^{-3} \text{ A}}{\frac{6.022 \cdot 10^{23}}{12 \cdot 10^{-6} \text{ m}^3/\text{mol}}} \cdot 1 \cdot 1.602 \cdot 10^{-19} \text{ As} \cdot 10 \cdot 10^{-6} \text{ m}^2$$

From:

<https://wiki.mexle.te.hs-heilbronn.de/> - MEXLE Wiki

Permanent link:

https://wiki.mexle.te.hs-heilbronn.de/ee2/task_tx86fewvysrcy8fc_with_calculation

Last update: 2024/07/05 00:14

