

task_ti7loik6aurfewkb_with_calculation

Student Group

First Name	Surname	Matrikel Nr.

Table of Contents

Exercise E1 Magnetic Flux Density (written test, approx. 6 % of a 120-minute written test,
SS2021) 2

magnetostatic, flux density, exam ee2 SS2021

Exercise E1 Magnetic Flux Density**(written test, approx. 6 % of a 120-minute written test, SS2021)**

A thin wire is placed horizontally in the laboratory. A current $I = 100 \text{ A}$ flows through it. The magnitude of $\hat{I} = 100 \text{ A}$ is operated.

Now stand next to it and think about what this value means. (3 points, independent)

The figure below shows the supply line between A and B and its position.

$$\text{Path } B = 0.12 \text{ m} \quad \mu_0 = 4\pi \cdot 10^{-7} \text{ Vs/Am}, \quad \mu_r = 1$$

The formula for the magnetic field strength can be rearranged:
$$H = \frac{I}{2\pi \cdot r} \quad r = \frac{I}{2\pi \cdot H}$$

Again, the magnetic flux density B is given as: $B = \mu_0 \cdot \mu_r \cdot H$

$$\text{Therefore: } r = \frac{\mu_0 \cdot \mu_r \cdot I}{2\pi \cdot B} \quad r = \frac{4\pi \cdot 10^{-7} \text{ Vs/Am} \cdot 100 \text{ A}}{2\pi \cdot 100 \cdot 10^{-6} \text{ T}}$$

a) What is the highest magnetic flux density through the line in your body? (3 points)

Path

The magnetic field strength for a conducting wire is given as:

$$\begin{aligned} H &= \frac{I}{2\pi \cdot r} \end{aligned}$$

The magnetic flux density B is given as: $B = \mu_0 \mu_r H$

Here, the maximum current is $\hat{I} = 100 \text{ ~\rm A}$ and the distance to the cable is $r = \sqrt{(0.1 \text{ ~\rm m})^2 + (0.4 \text{ ~\rm m})^2} = 0.412... \text{ ~\rm m}$.

Therefore:
$$B = 4\pi \cdot 10^{-7} \frac{\text{Vs}}{\text{Am}} \cdot 1 \cdot \frac{100 \text{ ~\rm A}}{2\pi \cdot 0.412... \text{ ~\rm m}}$$

From:

<https://wiki.mexle.te.hs-heilbronn.de/> - MEXLE Wiki

Permanent link:

https://wiki.mexle.te.hs-heilbronn.de/ee2/task_ti7loik6aurfewkb_with_calculation?rev=1719835260

Last update: **2024/07/01 14:01**

